

HYPERSPECTRAL MAPPING OF EARTH'S EARLIEST HYDROTHERMAL ACTIVITY IN AN ARCHEAN GRANITE-GREENSTONE TERRANE

Adrian J. Brown¹

1 Introduction

As part of a joint CSIRO-Australian Centre for Astrobiology (ACA) project to map hydrothermally altered minerals in the East Pilbara Granite-Greenstone Terrane (EPGGT), mineral maps of epithermal activity in two regions of volcanic successions have been compiled using an airborne hyperspectral dataset. The differing mineral suites suggest epithermal activity characterized by high and low sulphidation, possibly at the Archean seafloor or in a shallow marine environment.




Figure 1 – Location map for the North Pole Dome and overall dataset. Central Lat and long of the dataset are 119°E 21°S. Dataset is approximately 27km wide and north is up.

1.1 Motivation

In 2005, NASA will launch a hyperspectral VNIR-SWIR-IR instrument called the Compact Reconnaissance Infrared Spectrometer for Mars (CRISM) onboard the Mars Reconnaissance Orbiter (MRO) mission. In order to increase the probability of this mission encountering a Martian geobiological event of interest, this project is studying

¹ Australian Centre for Astrobiology, Macquarie University, NSW 2109, Australia.
corresponding email: abrown@els.mq.edu.au
website: <http://aca.mq.edu.au/abrown.htm>

some of the oldest, best preserved volcanic successions with life signs on Earth at the Pilbara Craton, Western Australia. Three water-related signs of interest exist in the target terrain:

- a. pillow basalts,
- b. stromatolites, and
- c. serpentinitised ultramafics.

The discovery of any of these water-signs on Mars would be of great significance for the understanding of the geological development of the planet and verifying the existence of an extant or extinct Martian biosphere.

2 Geological Setting

The study region discussed herein lies in a region called the North Pole Dome (NPD), a succession of ovoid shaped volcanic units surrounding a central monzogranite, interpreted as a syn-volcanic laccolith (Van Kranendonk, 2000). The volcanic units form part of the 3.5-3.3 Ga Warrawoona Group, and include ultrabasic komatiite, mafic and felsic units (Brown *et al.*, 2004a) that dip 30-70 degree away from the central monzogranite. A summary of the stratigraphy of the Warrawoona Group, along with associated fossil and stromatolite assemblages, is given in Table 1.

Age (Ga)	Unit	Fossil assemblages
3.458	Euro Basalt Strelley Pool Chert Panorama Formation	Well preserved conical stromatolites (Hofmann <i>et al.</i> , 1999)
3.470	Apex Basalt Duffer Formation Dresser Formation	Microfossils within Apex Chert (Schopf, 1993) Domical stromatolites and microfossils (Walter et al. 1980; Awramik 1983)
3.515	Mt Ada Basalt Coonterunah Group	

Table 1 – Stratigraphic column of Warrawoona Group units present at the North Pole Dome (Van Kranendonk, 2000). Dates are derived from U-Pb isotopes from zircons in the units and are accurate to approximately 3 million years from Thorpe (1992).

2.1 Hydrothermal Alteration

Submerged hydrothermal vents have been posited as a possible setting for the genesis of life (Nisbet and Sleep, 2001). In the search for signs of life in early Archean terrains, such as the Pilbara, hydrothermally altered zones are an attractive location to commence looking. In addition to the ability of hydrothermal activity to sustain life, it also has the capability of preserving life signs (Walter and Des Marais, 1993). These principles would be true for life on Mars, as they are for life here on the early Earth.

2.1.1 Dresser Formation

There are two primary hydrothermal events in evidence at the North Pole Dome. The first event, terminating at the Dresser Formation, is characterized by sub vertical black cherts leading up to a palaeo-surface, now replaced by chert, at the top of the Dresser Formation. The black cherts were first proposed as sedimentary features (Dunlop and Buick, 1981) but have now been interpreted as hydrothermal conduits (Nijman *et al.*, 2001), and carbon isotope geochemistry has lead some researchers to propose them as an early niche for life (Ueno *et al.*, 2004). Barite accompanies the black chert throughout the unit, and has been proposed as a syn-depositional exhalite product of the hydrothermal system (Van Kranendonk and Pirajno, in press). Domical stromatolites have been found at the extremities of Dresser Formation. This has led to some researchers suggesting the Dresser Formation was an early habitat of life (Groves *et al.*, 1981).

The style of alteration at the Dresser Formation is quite distinct from that found at a later horizon within the NPD, at the Strelley Pool Chert (SPC) level.

2.1.2 Strelley Pool Chert

The 3.2 Ga Strelley Pool Chert is a stromatolitic marker chert unit found in several regions throughout the Eastern Pilbara (Lowe, 1983). It outcrops distinctively at the outer extremities of the NPD, on top of the felsic Panorama Formation. It consists of carbonate and silicified stromatolites, black and white injection cherts and a succession of silicified clastic horizons. It is overlain by the pillow basalts of the Euro Basalt unit.

The presence of stromatolites (Hofmann *et al.*, 1999) and black and white injection cherts has lead some to propose a hydrothermal-based biosphere in the Early Archaean. This has been disputed (Lindsay *et al.*, 2003) but is an intriguing possibility. The underlying Panorama Formation is in places altered pervasively to pyrophyllite, and several sericite rich veins lead up to the SPC level, cutting vertically through the underlying Apex and Panorama Formations. These will be discussed further below.

3 Methods

A VNIR-SWIR hyperspectral dataset covering 600 sq. km of the NPD region was collected on 22 October 2002. The dataset was collected with the HyMap instrument (Cocks *et al.*, 1998), which is analagous to the AVIRIS instrument. The Pilbara coverage was collected as 14 swathes, each 2km wide. The instrument was flown at approximately 2.5km, or 8200ft AMSL. Spectral coverage was between 450-2500 nanometers in 126 contiguous bands. The date of the collection was timed to coincide with the end of the local dry season, in order to minimize vegetation coverage. The spatial resolution of the dataset is approximately 5m per pixel along track.

Mineral maps were produced using the continuum removed SWIR region by applying the “spectral feature fitting” method in the program ‘ENVI’ (www.rsi.com). Representative spectra were identified using a combination of spectral library comparison and ground testing through fieldwork. Hand samples were collected with the assistance of a PDA and GPS unit and were analysed using PIMA (hand held spectrometer), petrographic thin section, XRD and XRF.

4 Results

The dataset was delivered as radiance at sensor and was subsequently processed using the CSIRO developed ‘Hycorr’ program, which is based upon the ATREM program, to give approximate surface reflectance. In addition, using the software program ‘ENVI’, the dataset was treated by a ‘continuum removal’ algorithm (Clark *et al.*, 1987). This enhanced the features in the 2.0-2.4 micron SWIR region which are of most interest when detecting hydrothermally altered minerals.

Figure 2 displays representative continuum removed spectra of five different hydrothermal alteration minerals extracted from the HyMap dataset. Note the spectra appear similar in regions of the spectrum up to 2.0 microns. In the SWIR region, significant differences exist that enable them to be differentiated from each other.




Figure 2 – Continuum removed spectra of representative hydrothermal minerals extracted from the North Pole dataset. Each mineral is offset for clarity. Note differences between minerals are apparent in the 2.0-2.4 micron region.

Contrasting mineral assemblages were found at the Dresser Formation and Strelley Pool Chert levels, representative mineral maps are shown in Figure 3 and 4 below.

4.1 Dresser Formation

At the Dresser Formation level, it was possible to discern an intense zone of sericitic alteration capped by a thin serpentinitic alteration layer, overlain by an intense chlorite layer. The region is also cut by several late quartz porphyry veins unrelated to alteration at this level. These are rich in sericite and are seen to cut through all horizons. From field mapping, it was determined that the intense sericitisation was most likely due to pervasive and intense weathering of country rock basalts most likely affected through hydrothermal veins, now present as prominent black chert veins that cut through and terminate at a cap chert below the serpentine rich horizon. The serpentine rich layer was determined to be an ultramafic peridotite, most probably representing the basal layer of a komatiite succession (Brown *et al.*, 2004a). The overlying chlorite layer most probably represents seafloor alteration of the constituent basalts.

In addition to mapping the alteration minerals, black chert units that have been posited as feeder conduits for the Dresser Formation hydrothermal activity (Nijman *et al.*, 1998) can be seen clearly in the visible region of the spectrum, often due to thin Fe rich veneers that preferentially coat the chert. These are visible as east-west trending lineaments in Figure 3.

The hydrothermal activity that produced the sericite alteration most probably occurred in a low pH environment. The presence of large barite deposits within and surrounding the vein testifies to an oxidising fluid, if one is to accept the Archean environment was generally reducing (Ohmoto, 1997; Holland, 1999). These locally oxidizing conditions may have been created by a primitive biosphere living in the hydrothermal vents.




Figure 3 – Mineral map for the Dresser Formation region with context map on left showing coverage. Sericite is orange, serpentine light green (vertical strip in middle) and chlorite is yellow. East-west trending lineaments are massive black cherts, interpreted as feeder dykes for hydrothermal sites at the Dresser level (see text).

4.2 Strelley Pool Chert Mineral Assemblages

At the Strelley Pool Chert level, horizontal mineral assemblages of carbonate-chlorite are overlain by a thin but prominent sericite-pyrophyllite layer (at the Panorama felsic volcanic level) which is in turn overlain by a chlorite-rich layer. Sericite rich veins grade into pyrophyllite and then terminate at the Strelley Pool Chert layer. Some layers within the Strelley Pool Chert display sericite signatures (Brown *et al.*, 2004b) however there is no pyrophyllite within the Strelley Pool Chert.

It is hypothesized that this alteration style betrays the actions of an acid-sulfate hydrothermal system as described by previous researchers (White and Hedenquist, 1990). The presence of a thin continuous layer of pyrophyllite beneath a chert horizon and associated with sericite veins leading up to this horizon shows similarities with acid-sulfate Porphyry Cu argillic alteration, as portrayed in Figure 4 below (Bonham Jr. and Giles, 1983).




Figure 4 – Hydrothermal alteration at the Strelley Pool Chert level. Image on left shows coverage of central image. In central image, orange represents sericite, yellow chlorite and blue pyrophyllite. Image on right from (Bonham Jr. and Giles, 1983) showing typical mineral assemblages of an acid-sulfate epithermal vein.

5 Conclusion

This study into hydrothermal alteration amongst some of the best preserved, oldest rocks preserved on this planet, is continuing. Latest results are being published at the project website - <http://aca.mq.edu.au/abrown.htm>. Hyperspectral mapping has already demonstrated the ability to detect mineral assemblages that give vital clues to interpretation of the conditions within these hydrothermal systems.

In 2005, NASA will launch the hyperspectral VNIR-SWIR Compact Reconnaissance Infrared Spectrometer for Mars (CRISM) onboard the Mars Reconnaissance Orbiter (MRO). The instrument will have a spatial resolution of approximately 16m in high resolution mode (Murchie *et al.*, 2003). This will allow researchers to investigate signs of hydrothermal alteration using the critical SWIR region at scales approaching that expected for individual veins of an epithermal deposit. Application of the lessons learnt from mapping volcanic terrains such as the Warrawoona Group in Western Australia will be of vital importance in informing the search for possible past or present habitats of life on Mars.

6 Acknowledgements

This project was supported by field equipment from Geological Survey of Western Australia. Core funding was provided by CSIRO and Macquarie University. Prof. Malcolm Walter and Dr. Thomas Cudahy are thanked for their continually inspirational supervision.

7 References

- Bonham Jr., H. F. and Giles, D. L. (1983) Epithermal gold/silver deposits - the geothermal connection, pp. 384, Geothermal Resources Council.
- Brown, A. J., Walter, M. R. and Cudahy, T. J. (2004a) Hyperspectral and field mapping of an Archaean Komatiite Unit in the Pilbara Craton, Western Australia: Applications for CRISM Mission, in *LPSC XXXV*, LPI, Houston.
- Brown, A. J., Walter, M. R. and Cudahy, T. J. (2004b) Remote Mapping of Earth's Earliest Biosphere: Hyperspectral Imagery of the Pilbara Craton and applications for finding niches for life on Mars, in *International Journal of Astrobiology*, pp. 16-17, Special Supplement 1 - Abstracts from the Astrobiology Science Conference 2004, NASA Ames Research Laboratory, Mountain View, CA.
- Clark, R. N., King, T. V. V. and Gorelick, N. (1987) Automatic Continuum Analysis of Reflectance Spectra, in *Proceedings of the Third Airborne Imaging Spectrometer Data Analysis Workshop*, pp. 138-142, JPL Publication 87-30, Pasadena, CA.
- Cocks, T., Janssen, R., Stewart, A., Wilson, I. and Shields, T. (1998) The HyMap Airborne Hyperspectral Sensor: The System, Calibration and Performance. *1st EARSeL Conference*, 37-42.
- Dunlop, J. S. R. and Buick, R., (1981) Archaean epiclastic sediments derived from mafic volcanics, North Pole, Pilbara Block, Western Australia, in *Special Publication of the Geological Society of Australia*, edited by D. I. Groves and J. E. Glover, pp. 225-233, Geological Society of Australia, Perth, WA.
- Groves, D. I., Dunlop, J. S. R. and Buick, R. (1981) An early habitat of life. *Scientific American*, 245 (4), 56-65.
- Hofmann, H. J., Grey, K., Hickman, A. H. and Thorpe, R. I. (1999) Origin of 3.45 Ga coniform stromatolites in Warrawoona Group, Western Australia. *GSA Bulletin*, 111 (8), 1256-1262.
- Holland, H. D. (1999) When did the Earth's atmosphere become oxic? A Reply. *The Geochemical News*, 100, 20-22.
- Lindsay, J. F., Brasier, M. D., McLoughlin, N., Green, O. R., Fogel, M., McNamara, K. M., Steele, A. and Mertzman, S. A. (2003) Abiotic Earth - Establishing a baseline for Earliest Life, Data from the Archean of Western Australia, in *LPSC XXXIV*, pp. 1137, LPI, Houston, TX.
- Lowe, D. R. (1983) Restricted shallow-water sedimentation of early Archean stromatolitic and evaporitic strata of the Strelley Pool Chert, Pilbara Block, Western Australia. *Precambrian Research*, 19 (3), 239-283.

- Murchie, S., Arvidson, R., Beisser, K., Bibring, J.-P., Bishop, J., Boldt, J., Bussey, B., Choo, T., Clancy, R. T., Darlington, E. H., Des Marais, D., Fasold, M., Fort, D. E., Green, R. O., Guinness, E. A., Hayes, J., Heyler, G., Humm, D., Lee, R., Lees, J., Lohr, D., Malaret, E., Morris, R. L., Mustard, J., Rhodes, E., Robinson, M., Roush, T., Schaefer, E., Seagrave, G., Silverglate, P., Smith, M. D., Strohbenn, K., Thompson, P. and Tossman, B. (2003) CRISM: Compact Reconnaissance Imaging Spectrometer for Mars on the Mars Reconnaissance Orbiter, in *Sixth International Conference on Mars*, pp. 3062, LPI, Houston, TX.
- Nijman, W., de Bruijne, K. H. and Valkering, M. E. (1998) Growth fault control of Early Archaean cherts, barite mounds and chert-barite veins, North Pole Dome, Eastern Pilbara, Western Australia. *Precambrian Research*, 88 (1-4), 25-52.
- Nijman, W., de Vries, S. T. and Houtzager, O. (2001) Earth's Earliest Sedimentary Basins: the Lower Archaean of the Pilbara and Kaapvaal compared, in *4th International Archaean Symposium*, edited by K. F. Cassidy; J. M. Dunphy and M. J. Van Kranendonk, pp. 520-522, AGSO, Perth, WA.
- Nisbet, E. G. and Sleep, N. H. (2001) The habitat and nature of early life. *Nature*, 409 (6823), 1083-1091.
- Ohmoto, H. (1997) When did the Earth's atmosphere become oxic? *The Geochemical News*, 93, 12-27.
- Ueno, Y., Yoshioka, H., Maruyama, S. and Isozaki, Y. (2004) Carbon isotopes and petrography of kerogens in ~3.5-Ga hydrothermal silica dikes in the North Pole area, Western Australia. *Geochimica et Cosmochimica Acta*, 68 (3), 573-589.
- Van Kranendonk, M. J. (2000) Geology of the North Shaw 1:100 000 Sheet, Geological Survey of Western Australia, Department of Minerals and Energy, Perth, WA.
- Van Kranendonk, M. J. and Pirajno, F. (in press) Geochemistry of metabasalts and hydrothermal alteration zones associated with ca. 3.45 Ga chert+/- barite deposits: implications for the geological setting of the Warrawoona Group, Pilbara Craton, Australia. *Geochemistry: Exploration, environment and analysis*.
- Walter, M. R. and Des Marais, D. J. (1993) Preservation of Biological Information in Thermal-Spring Deposits - Developing a Strategy for the Search for Fossil Life on Mars. *Icarus*, 101 (1), 129-143.
- White, N. C. and Hedenquist, J. W. (1990) Epithermal environments and styles of mineralization: variations and their causes, and guideline for exploration. *Journal of Geochemical Exploration*, 36, 445-474.